

Trichoderma cellulase isolated from a culture of Trichoderma longibrachiatum with an estimated molecular weight of 95-105 kD and an estimated pH optimum of 5.0 at 338 Kelvin and ph 4.0 at 318 and 328 Kelvin

Patent number: NZ335107
Publication date: 1999-11-29
Inventor: CLARKSON KATHLEEN A; COLLIER KATHERINE D; KELLIS JAMES T; KELLY MOIRA B; LARENAS EDMUND A; BOWER BENJAMIN S
Applicant: GENENCOR INT
Classification:
- International: A23K1/165; C11D3/386; C12N9/42; D06M16/00; D06P5/02; D06P5/15; D21C5/00; A23K1/165; C11D3/38; C12N9/42; D06M16/00; D06P5/02; D06P5/15; D21C5/00; (IPC1-7): C07K14/37; A23K1/165; C08B30/04; C11D3/386; C12N1/14; C12N9/42; C12N15/56; D06M16/00; D21C5/00
- european: A23K1/165B; C11D3/386F; C12N9/42; D06M16/00B; D06P5/02; D06P5/15E; D21C5/00B
Application number: NZ19970335107 19971007
Priority number(s): US19960728350 19961009; WO1997US18402 19971007

Also published as:

- WO9815619 (A1)
- EP0934402 (A1)
- US6017870 (A1)
- EP0934402 (B1)
- DE69726748T (T2)
- AU719674 (B2)

less <<

[Report a data error here](#)

Abstract of NZ335107

The present invention relates to a purified cellulase composition, where the purified cellulase composition corresponding to a naturally occurring cellulase produced during a fermentation of Trichoderma longibrachiatum which exhibits a molecular weight of about 95-105 kD as measured on SDS-PAGE gel and a pI of about 5.6-6.8 as measured on an IEF gel. Also claimed is a method for preparing a purified cellulase composition comprising the steps: a) preparing a fermentation culture of a microorganism which produces a cellulase composition corresponding to a naturally occurring cellulase produced during fermentation of Trichoderma longibrachiatum which exhibits a molecular weight of about 95-105 kD as measured on SDS-PAGE gel and a pI of about 5.6-6.8 as measured on an IEF gel and b) separating the cellulase from the microorganisms to form a purified cellulase composition

Data supplied from the **esp@cenet** database - Worldwide